



## REVIEW ARTICLE

# Cardiopulmonary exercise testing as a risk assessment method in non cardio-pulmonary surgery: a systematic review

T. B. Smith,<sup>1</sup> C. Stonell,<sup>2</sup> S. Purkayastha<sup>3</sup> and P. Paraskevas<sup>4</sup>

*1 Medical Student, 2 Consultant, Department of Anaesthesia, 3 Clinical Lecturer, 4 Senior Lecturer and Consultant Surgeon, Departments of Biosurgery and Surgical Technology and Academic Surgical Unit, St Mary's Hospital, Imperial College Healthcare NHS Trust, London, UK*

### Summary

This study reviews the predictive value of maximum oxygen consumption ( $\dot{V}O_{2\max}$ ) and anaerobic threshold, obtained through cardiopulmonary exercise testing, in calculating peri-operative morbidity and mortality in non-cardiopulmonary thoraco-abdominal surgery. A literature review provided nine studies that investigated either one or both of these two variables across a wide range of surgical procedures. Six of the seven studies that reported sufficiently detailed results on peak oxygen consumption and four of the six studies that reported sufficiently detailed results on anaerobic threshold found them to be significant predictors. We conclude that peak oxygen consumption and possibly anaerobic threshold are valid predictors of peri-operative morbidity and mortality in non-cardiopulmonary thoraco-abdominal surgery. These indicators could potentially provide a means of allocating increased care to high-risk patients.

Correspondence to: Dr Christopher Stonell

E-mail: christopher.stonell@imperial.nhs.uk

Accepted: 1 April 2009


Cardiopulmonary exercise testing (CPET or CPX) is a non-invasive, integrated assessment of cardiovascular and pulmonary function both at rest and under stress. Among the purported benefits is its ability to determine the ability of the subject's physiological capacity to cope with the metabolic demands created by the trauma of major surgery. During the test subjects are exposed to incremental physical exercise up to their maximally tolerated level, dictated either by exhaustion or symptom related cessation (e.g., breathlessness or angina). Several physiological variables are recorded including ventilatory parameters, inspiratory and expiratory gases, blood pressure (BP) and electrocardiogram (ECG). From these are derived two key indicators: the body's maximum oxygen uptake ( $\dot{V}O_{2\max}$ ) and the point at which anaerobic metabolism exceeds aerobic metabolism (Ventilatory Anaerobic Threshold or VAT). Together, these broadly indicate the ability of the cardiovascular system to deliver oxygen to the peripheral tissues and the ability of the tissues to utilise that oxygen. It has already been demonstrated that measures such as  $\dot{V}O_{2\max}$  are

useful predictors of postoperative complications of pulmonary resection surgery [1–3] and assessing the timing of cardiac transplant surgery [4], while the VAT is a predictor of postoperative cardiac complications in abdominal surgery [5].

### Physiology of the anaerobic threshold


Lactate accumulation in exercising muscle occurs when the muscle  $O_2$  demand exceeds the supply. At this point the venous lactate concentration will begin to rise. This point is called the Lactate Anaerobic Threshold (LAT). The muscle lactate/pyruvate ratio increases at the LAT which supports the concept that the lactic acidosis results from relative muscle hypoxia [6–8].

As an individual commences an incremental exercise test, their expired minute volume ( $\dot{V}_E$ ), oxygen consumption per minute ( $\dot{V}O_2$ ) and  $CO_2$  production per minute ( $\dot{V}CO_2$ ) all increase linearly with respect to variables such as work rate or time. However a point is reached when  $\dot{V}CO_2$  increases out of proportion to  $\dot{V}O_2$




**Figure 1**  $\dot{V}O_2$  (---) and  $\dot{V}CO_2$  (·····) vs work rate:  $\dot{V}O_2$  and  $\dot{V}CO_2$  initially increase linearly and proportionately to the work rate in early exercise. A point is reached where  $\dot{V}CO_2$  increases and exceeds  $\dot{V}O_2$  (arrow). Towards maximum exercise  $\dot{V}O_2$  reaches a plateau while  $\dot{V}CO_2$  continues to rise.

(Fig. 1). This change is attributed to  $HCO_3^-$  buffering the lactate produced and consequently generating a relative excess of  $CO_2$ . This transition is termed the Lactate Anaerobic Threshold (LAT). This phenomenon allows determination of the anaerobic threshold by respiratory measurements. Generally this is seen at 47–64% of  $\dot{V}O_{2max}$  in healthy untrained individuals [9] and is a marker of the maximum work rate that can be sustained for a prolonged period [10, 11]. As this point is less effort dependant than  $\dot{V}O_{2max}$  it is a clinically attractive measurement in a patient population where it may be difficult to achieve the maximal effort required for a true  $\dot{V}O_{2max}$  result. Failure to perform sufficient exercise during the test to reach VAT suggests either poor motivation or a non-cardiac (e.g. pulmonary or musculoskeletal) problem [12]. On reaching the AT a fall in the arterial pH caused by anaerobic respiration results in the carotid bodies stimulating an increase in minute ventilation [13]. This point is termed the Ventilatory Anaerobic Threshold (VAT) (Fig. 2). During this period the  $\dot{V}_E$  increases in proportion to the  $\dot{V}CO_2$  and the  $\dot{V}_E/\dot{V}CO_2$  remains constant. The rate of rise of  $\dot{V}_E$  exceeds the rate of rise of  $\dot{V}O_2$  and so  $\dot{V}_E/\dot{V}O_2$  starts to increase. This gives one method of determining the VAT from gas exchange measurements as AT is the point when the Ventilatory Equivalent for oxygen ( $\dot{V}_E/\dot{V}O_2$ ) increases relative to a constant Ventilatory Equivalent for carbon dioxide ( $\dot{V}_E/\dot{V}CO_2$ ) [14] (Fig. 3).




**Figure 2** Expired minute ventilation vs work rate. The VAT occurs when the  $\dot{V}_E$  slope begins to steepen (arrow) during an incremental exercise test, indicating an increase in  $\dot{V}_E$  due to carotid body stimulation as a result of lactate accumulation.



**Figure 3** Determining AT by ventilatory equivalents of  $O_2$  ( $\dot{V}_E/\dot{V}O_2$ ) and  $CO_2$  ( $\dot{V}_E/\dot{V}CO_2$ ): with incremental work the  $\dot{V}_E/\dot{V}O_2$  (·····) and  $\dot{V}_E/\dot{V}CO_2$  (—) initially decrease to reach a plateau. The AT is determined by marking the point (arrow) at which  $\dot{V}_E/\dot{V}O_2$  starts to increase while  $\dot{V}_E/\dot{V}CO_2$  remains constant or falls slightly (due to  $\dot{V}_E$  increasing disproportionately to  $\dot{V}O_2$  but proportionately to  $\dot{V}CO_2$ ). The work rate at which AT occurs can then be derived (arrow).

Another common method used to determine VAT is the ‘V-Slope’ method [15]. In order to determine the VAT  $\dot{V}CO_2$  is plotted against  $\dot{V}O_2$  during an incremental exercise test. Initially the slope is unitary, however at the VAT, when  $\dot{V}CO_2$  begins to increase, the slope increases. From this point of deflection of  $\dot{V}CO_2$  the  $\dot{V}O_2$  at which AT is reached can be determined (Fig. 4). Thus there



**Figure 4** V-slope method of determining AT:  $\dot{V}CO_2$  is plotted against  $\dot{V}O_2$  (bold line). Using the method described by Beaver et al. [15], linear regression lines are drawn through the lower and upper parts of the curve. The inflection point corresponds to the intersection of the two lines (arrow). This indicates the  $\dot{V}O_2$  at which the AT occurs. This is essentially the same data as Fig. 3 plotted in a different way in order to determine the transition between aerobic and anaerobic metabolism.

are several methods, using measures of physiological responses to lactate production, which can be employed to non-invasively estimate the transition from predominantly aerobic metabolism to predominantly anaerobic metabolism during exercise.

Whether lactate appears suddenly or if there is gradual accumulation over a period of time is debated [16–21]. If this former hypothesis were correct it would make the VAT less relevant (in contrast to LAT) in providing an accurate indication of the onset of anaerobic respiration. Furthermore, there is some evidence to suggest that LAT and VAT may not be directly linked. Paterson et al. [22] investigated patients with McArdle's disease (a myophosphorylase deficiency which makes subjects unable to produce lactic acid during exercise) and found a similar non-linear ventilatory response to exercise as seen in normal subjects. Additionally studies using dopamine infusion [23] or high oxygen [24] to inhibit carotid body function have demonstrated persistence of a VAT.

Cardiopulmonary exercise testing relies upon accurate breath-by-breath measurements of pulmonary gas exchange at the mouth to represent transfer at the pulmonary capillaries. However in practice they are not identical and so there are several models that propose to estimate the transfer at the pulmonary capillaries from values recorded at the mouth [25, 26]. These models produce different levels of breath-to-breath variation in gas exchange and it is not yet understood whether this variability is a by-product of the conversion method or an actual physiological occurrence [27]. This has potential to add a degree of error to accurately assigning points such as

VAT and would benefit from further elucidation on the subject.

Notwithstanding these important issues needed to clarify the physiology underlying VAT and LAT, it is still possible to use either or both as a (perhaps relatively crude) non-invasive and useful marker of the onset of a significant anaerobic component in respiration.

### Cardiopulmonary exercise testing in practice

There are several different protocols for achieving  $\dot{V}O_{2\max}$ . Most use either an exercise bicycle (cycle ergometer) or a treadmill. Cycle ergometers tend to be used in peri-operative clinical settings as they provide a degree of safety in maximal testing, offset the work of walking in the obese and are less prone to artifacts in the ECG, airflow and pressure measurement. A higher peak  $\dot{V}O_2$  can usually be generated on a treadmill [9, 12], however there is a learned technique to walking on a treadmill, which patients may not be familiar with, and should the subject hold on to any part of the treadmill the amount of work performed is altered as a consequence of static work being performed in the arms. This would interfere with the determination of an accurate  $\dot{V}O_2$ -work rate relationship as the static work performed is not accounted for by the ergometer [9]. In both methods, work rate is gradually and imperceptibly increased in a stepped or ramped manner until the subject is unable to continue. On a treadmill this can be achieved by increasing both the gradient and speed while on an exercise bicycle it is achieved by increasing the resistance of the pedals while the subject maintains a constant pedaling rate. If a subject is unable to cycle or walk then an arm crank may be employed.

The optimal duration of exercise is approximately 10 min [28]. Too short a duration of exercise may produce an underestimation of  $\dot{V}O_{2\max}$  [28], while too long a duration may cause premature cessation due to the subject becoming demotivated or because of orthopaedic factors [29]. VAT seems to be independent of the duration of exercise [28]. The testing protocol can be adjusted according to predicted values based on age and height to achieve this [30]. In the United Kingdom the Consensus Protocol for Pre-Operative CPX Testing for York, Torbay and UCLH (Pre-operative Association, UK) is an accepted pre-operative testing protocol.

All physiological variables are recorded breath by breath and respiratory variables are then typically averaged over periods of 15–45 s [29].

The term  $\dot{V}O_{2\max}$  is used for the highest  $\dot{V}O_2$  recorded when a subject's  $\dot{V}O_2$  is observed to reach a plateau with increasing workload. If this plateau is not seen, as is often

the case, but the subject is believed to be near to their maximal effort it is termed their  $\dot{V}O_{2\text{peak}}$ . In a clinical setting these are considered essentially similar.

### Current applications of CPET

The American Thoracic Society/American College of Chest Physicians Statement on Cardiopulmonary Exercise Testing (2003) lists the specific indications for CPET testing [31] that include:

- 1 Evaluation of exercise tolerance, where the diagnosis is known, in order to objectively evaluate functional capacity, disability or response to treatment.
- 2 Evaluation of undiagnosed exercise intolerance where cardiac and respiratory aetiologies may coexist, the symptoms are disproportionate to the results of resting investigations or the investigations are non diagnostic.
- 3 Evaluation of patients with cardiovascular diseases.
- 4 Evaluation of patients with respiratory diseases/symptoms.
- 5 Pre-operative evaluation.
- 6 Exercise evaluation and prescription for pulmonary rehabilitation.
- 7 Evaluation of impairment/disability.
- 8 Evaluation for lung, heart and heart-lung transplantation.

We explain some of these in greater detail:

*Evaluation of exercise tolerance:* As  $\dot{V}O_{2\text{max}}$  measures maximal aerobic capacity it is a very useful indicator of the 'cardiovascular fitness' of an athlete, an important variable in endurance sports that have a high aerobic demand such as long distance running, cycling and skiing. However as most athletes at a specific level of a sport will have similar values it does not serve to predict their individual competitive performance (as other variables such as economy come into play) but serves instead to act as a personal indicator after a period of training or injury through comparison with previous values (e.g., akin to a golf handicap).

*Evaluation of impairment/disability:* Oxygen consumption can be used to indirectly assess caloric expenditure in subjects, since the two mirror each other during exercise. This results in the calculation of a Metabolic Equivalent (MET), with 1 MET corresponding to an oxygen usage of  $3.5 \text{ ml.kg}^{-1}.\text{min}^{-1}$  (or the same resting intake as a healthy 70 kg man) [29]. It is therefore possible to establish the level of physical activity a subject should be able to obtain in reference to common activities. For example making a bed has a metabolic cost of 2 MET's, painting a fence 4.5 MET's, jogging 7 MET's and moving furniture 9 METS [32]. With this reference data and the known  $\dot{V}O_{2\text{max}}$  of an individual it is possible to determine those daily tasks that can be completed

independently and those for which assistance is needed.  $\dot{V}O_{2\text{max}}$  rather than VAT is more accurate in this context since individuals with a very low  $\dot{V}O_{2\text{max}}$  may be producing lactate at rest [33].

### Cardiopulmonary testing as a pre-operative predictor of risk

Cardiopulmonary exercise testing is most documented in cardiac and pulmonary medicine and surgery. A review of multiple lung tumour resection studies by Beckles et al. [34] provides data to suggest  $\dot{V}O_{2\text{max}}$  can be used to predict peri-operative complication rates in tumour resection surgery:

- 1  $> 20 \text{ ml.kg}^{-1}.\text{min}^{-1}$  = no increased risk of complications or death,
- 2  $< 15 \text{ ml.kg}^{-1}.\text{min}^{-1}$  = increased risk of peri-operative complications,
- 3  $< 10 \text{ ml.kg}^{-1}.\text{min}^{-1}$  = a very high risk of for post-operative complications.

Mancini et al. [4] used CPET to assess the urgency of cardiac transplantation by comparing the survival of heart failure patients who were not accepted for cardiac transplantation vs those who received a transplant. Of those not accepted for transplantation, a  $\dot{V}O_{2\text{peak}} > 14 \text{ ml.kg}^{-1}.\text{min}^{-1}$  yielded survival comparable to those after transplantation; a  $\dot{V}O_{2\text{peak}} < 10 \text{ ml.kg}^{-1}.\text{min}^{-1}$  yielded lower survival.

However few studies have investigated the role of cardiopulmonary exercise testing in the pre-operative assessment of patients undergoing non-cardiopulmonary thoraco-abdominal surgery. If CPET was able to provide a reliable indicator of peri-operative morbidity and mortality it could have a significant impact on the care of surgical patients more generally and on the cost of their healthcare. By identifying high risk candidates additional resources could be put in place ahead of time (e.g. an Intensive Care bed) to ensure appropriate postoperative care and thus potentially reduce morbidity and mortality. Furthermore if candidates identified as being at risk were placed on an exercise regime before elective surgery this could potentially improve their CPET measurements, and in turn their postoperative risk.

In cardiac surgery, and some forms of pulmonary surgery, it is expected that cardiac and respiratory function will improve postoperatively. In contrast, after non-cardiothoracic surgery cardio-respiratory function will remain, at best, unaffected and may well deteriorate postoperatively, at a time that the demands placed on the cardio-respiratory system are increased. We used the method of systematic review to establish the role of CPET testing in non-cardiopulmonary surgery.

## Methods of systematic review

### Study selection

Pubmed and ISI Web of Knowledge were searched for studies investigating pre-operative exercise testing as an indicator of postoperative outcome in surgery. The search terms used were ‘CPET/surgery’, ‘CPEX/surgery’, ‘cardiopulmonary/exercise testing/surgery’ and ‘ $\dot{V}O_{2max}$ /surgery’. With both search engines the ‘related articles’ or ‘related records’ function was used to broaden the search. The references of the studies of interest were reviewed as were any studies which were found to cite them.

### Inclusion/exclusion criteria

In order to be included studies had to compare subjects pre-operative  $\dot{V}O_{2max}$  and/or AT with their postoperative outcome in terms of morbidity or mortality. Studies were excluded if they reported on (i) pulmonary or cardiac surgery, (ii) non-thoraco-abdominal surgery. The selected studies were then analysed for data investigating the relationship between pre-operative CPET values, such as  $\dot{V}O_{2max}$  or AT, to postoperative outcomes such as mortality or a defined morbidity. Where present this data was extracted and included in the review.

## Results

In all, 322 studies were initially identified, 315 of which were excluded (Fig. 5). The reference lists of the seven remaining studies yielded two additional articles (Table 1).

All but one study was in English [39] which was written in Japanese with an English abstract. The abstract was deemed to have sufficient data for the purposes of this review. The nine studies can be divided into




Figure 5 Profile of literature review.

four categories: abdominal aortic aneurysm repair (AAA;  $n = 2$ ), hepatic transplantation ( $n = 1$ ), upper GI surgery ( $n = 4$ ) and major intra-abdominal surgery in the elderly ( $n = 2$ ).

### Abdominal aortic aneurysm repair

Carlisle et al. [35] measured AT,  $\dot{V}O_{2peak}$ ,  $\dot{V}_E/\dot{V}O_2$  and  $\dot{V}_E/\dot{V}CO_2$  in 130 patients before elective repair of an AAA. All variables correlated with survival (median follow up 35 months) in univariate analysis;  $\dot{V}O_{2peak}$  Hazard ratio (95% CI [standard error of the hazard ratio (SE)]) was 0.83 (0.74–0.94 [0.05])  $p = 0.002$  and AT 0.74 (0.63–0.87 [0.06])  $p < 0.001$ . In multivariate analysis however  $\dot{V}_E/\dot{V}CO_2$  correlated best with survival to both 30 days and the total observation period 1.13 (1.07–1.19 [0.03])  $p < 0.001$ . AT was also associated with midterm survival but to a lesser degree 0.84 (0.72–0.98 [0.07])  $p = 0.033$ ,  $\dot{V}O_{2peak}$  was not.

Nugent et al. [41] recorded the  $\dot{V}O_{2peak}$  of 30 patients prior to elective AAA repair and followed them up postoperatively for 12 months. He found no significant difference in the  $\dot{V}O_{2peak}$  in those who had postoperative complications compared to those who did not, though there was an underlying trend with 70% of the patients with complications having a  $\dot{V}O_{2peak}$  of  $< 20 \text{ ml}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$  as opposed to 50% in the non-complication group. However this study did not specify any power calculations for their analysis and compared to other studies in the review had the smallest number of subjects. Therefore findings from this group should be treated with caution.

These two studies suggest that there is a weak relationship between  $\dot{V}O_{2peak}$  and postoperative morbidity and mortality in patients undergoing elective AAA repair. Only one study investigates the role of AT but finds it to be a stronger predictor.

### Hepatic transplantation

Fifty nine subjects who underwent hepatic transplantation [36] and were followed up 100 days postoperatively, were subsequently divided into two groups, survivors ( $n = 53$ ) and non-survivors ( $n = 6$ ). Non-survivors were more likely to have a peak  $\dot{V}O_2 < 60\%$  predicted ( $p = 0.04$ ) and a  $\dot{V}O_2\text{-AT} < 50\%$  predicted ( $p = 0.03$ ). They were also more likely to have a combined peak  $\dot{V}O_2 < 60\%$  predicted and a  $\dot{V}O_2\text{-AT} < 50\%$  predicted ( $p < 0.01$ ). This study thus suggests that both  $\dot{V}O_{2peak}$  and AT provide an indication of short term mortality following hepatic transplantation.

### Upper GI surgery

Seventy-eight patients underwent oesophagectomy [37], and comparison between patients who did or did not



Table 1 Study characteristics.

Author (References)	Type of surgery	Year	Country of origin	Design	Total patients
Carlisle et al. [35]	AAA repair	2007	UK	Bicycle	130
Epstein et al. [36]	Hepatic transplantation	2004	USA	Bicycle	59
Forshaw et al. [37]	Oesophagectomy	2008	UK	Bicycle	78
McCullough et al. [38]	Roux-en-Y gastric bypass	2006	USA	Treadmill	109
Nagamatsu et al. [39]	Thoraco-lapartomy for oesophageal cancer	1994	Japan	Not specified	52
Nagamatsu et al. [40]	Oesophagectomy and lymphadenectomy.	2001	Japan	Bicycle	91
Nugent et al. [41]	AAA repair	1998	Northern Ireland	Treadmill	36
Older et al. [42]	Elderly (intra-abdominal surgery)	1993	Australia	Bicycle	187
Older et al. [5]	Elderly (intra-abdominal surgery)	1999	Australia	Bicycle	548

experience postoperative cardiopulmonary complications showed significant difference in their  $\dot{V}O_{2\text{peak}}$  ( $p = 0.04$ ) and although there was a trend in their AT, this was not significant ( $p = 0.07$ ).  $\dot{V}O_{2\text{peak}}$  and AT were also analysed by receiver operating characteristic curves with values of 0.63 (95% CI 0.50–0.76) and 0.62 (95% CI 0.49–0.75) respectively. Although the authors state that these values were significant different from 0.5 ( $p = 0.03$  and 0.02 respectively) the values are under 0.65 and so do not show strong predictive ability.

In the earlier of Nagamatsu et al.'s [39] two papers, subjects were divided into two groups depending on the presence ( $n = 11$ ) or absence ( $n = 41$ ) of cardiopulmonary complications. There were significant differences between the two groups in their pre-operative  $\dot{V}O_{2\text{max}} \cdot \text{m}^{-2}$  ( $p < 0.001$ ) and  $\text{AT} \cdot \text{m}^{-2}$  ( $p < 0.001$ ).

Nagamatsu et al.'s [40] later paper investigated CPET in 91 patients undergoing radical oesophagectomy with three-field lymphadenectomy. Subjects were divided into two groups postoperatively based on cardiopulmonary complications. It was observed that  $\dot{V}O_{2\text{max}} \cdot \text{m}^{-2}$  was lower among patients having cardiopulmonary complications ( $p < 0.001$ ), though this was not seen for  $\text{AT} \cdot \text{m}^{-2}$  values. Subjects were also grouped postoperatively into one of seven groups by their  $\dot{V}O_{2\text{max}} \cdot \text{m}^{-2}$  and the cardiopulmonary complications for each group calculated. This was 86% with a  $\dot{V}O_{2\text{max}} \cdot \text{m}^{-2} < 699 \text{ ml} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$ , 44% with a  $\dot{V}O_{2\text{max}} \cdot \text{m}^{-2}$  of 700–799  $\text{ml} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$ , 10% with a  $\dot{V}O_{2\text{max}} \cdot \text{m}^{-2}$  of 800–1099  $\text{ml} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$  and 0% with a  $\dot{V}O_{2\text{max}} \cdot \text{m}^{-2}$  of 1100  $\text{ml} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$  or more.

McCullough et al. [38] recorded the  $\dot{V}O_{2\text{peak}}$  and AT of 109 bariatric patients (i.e., body mass index (SD) 48.7 (7.2)) prior to laparoscopic Roux-en-Y gastric bypass surgery. Patients were divided into tertiles based on their  $\dot{V}O_{2\text{peak}}$  (1: 6.8–15.8  $\text{ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ , 2: 15.9–18.4  $\text{ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  and 3: 18.5–27.7  $\text{ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ) and their complication rates were compared between these groups. It was found that complications occurred in 16.6% of patients with a peak  $\dot{V}O_2 < 15.8 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  and in only 2.8% of patients with a peak  $\dot{V}O_2 > 15.8 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,

univariate odds ratio 6.77 (95% CI 1.14–51.76)  $p = 0.02$ . Multivariate analysis using peak  $\dot{V}O_2$  as a continuous variable showed it was a significant predictor of complications, odds ratio of 1.61 per unit decrease (95% CI 1.19–2.18)  $p = 0.002$ . Hospital length of stay and 30-day readmission rates were highest in the lowest tertile, 3.8 days vs 2.8 days for all others ( $p = 0.002$ ). The AT was not investigated as a postoperative indicator but showed a significant trend across the tertiles ( $p < 0.001$ ).

The first three studies all show that patients who experience cardiopulmonary complications after oesophagectomy, as a group, have significantly different  $\dot{V}O_{2\text{peak}}$  from patients who do not experience cardiopulmonary complications. Nagamatsu et al.'s [40] later study chose to use 800  $\text{ml} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$  as the minimal acceptable value for patients undergoing oesophagectomy. However only one of the three papers [39] saw a significant difference in AT between the complication and non-complication groups. The final study on bariatric patients [38] shows that their  $\dot{V}O_{2\text{peak}}$  played a predictive role on their postoperative complication rate, however this was only between the lowest tertile and the two higher ones.

### Intra-abdominal surgery in the elderly

Older et al. [42] recorded the pre-surgical anaerobic threshold of 187 elderly patients undergoing major abdominal surgery (abdominal surgery which would be 'likely to cause a significantly increase in oxygen demand' [42]). It was found that in patients with an AT of  $< 11 \text{ ml} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$  ( $n = 55$ ) 10 died of cardiovascular causes (18%), while in patients with an AT of  $> 11 \text{ ml} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$  ( $n = 132$ ) only one died of cardiovascular causes (0.8%),  $p < 0.001$ . When a low AT was combined with known pre-operative ischaemia ( $n = 19$ ) the mortality rate was 42%, as opposed to 4% in those with a high AT and pre-operative ischaemia ( $n = 25$ ). This established that AT and ischemia were important risk factors for surgery in the elderly and led the

authors to suggest that all patients scheduled for major intra-abdominal surgery with an AT  $< 11 \text{ ml}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$  should be admitted to ICU pre-operatively. Older et al. [5] then subsequently investigated the impact of triaging patients, based on this data, on their postoperative outcome.

In this latter study [5] 548 patients over the age of 60 (or younger if they had cardiopulmonary disease) were assigned into one of three groups pre-operatively: ICU, HDU or ward admission. This was based on their pre-operative AT and ECG ST segment changes, which were ascertained by CPET. If a patient had an AT  $< 11 \text{ ml}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$  they were assigned to ICU pre-operatively (additionally if a patient was undergoing a high risk surgery e.g. oesophagectomy or aortic they were also assigned to this group). This accounted for 28% of the patients. If a patient had an AT  $> 11 \text{ ml}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$  but either demonstrated myocardial ischemia (on CPET) or a  $\dot{V}_E/\dot{V}O_2$  of  $> 35$  they were admitted to HDU post-operatively (21%). All other patients were sent to the general ward after surgery (51%). Of the nine patients who died postoperatively from cardiopulmonary complications, seven had an AT  $< 11 \text{ ml}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$  and the other two were in the HDU category. There were no deaths in the ward admissions from cardiopulmonary causes.

These two papers show that AT has a predictive value in determining postoperative complications across a range of surgical procedures in the elderly.

### Results summary

Seven of the nine studies reviewed sufficiently investigated the predictive value of pre-operative  $\dot{V}O_{2\text{max/peak}}$  for postoperative complications, of these six found  $\dot{V}O_{2\text{max/peak}}$  to be a valid indicator. Fewer studies reported investigating the role of Anaerobic Threshold in this regard but in the six that did so adequately, four also found it to be a valid indicator (see Table 2).

### Discussion

This review demonstrates that the majority of investigations of CPET as an indicator of peri-operative complications, in non-cardiopulmonary surgery, find the technique, and  $\dot{V}O_{2\text{peak}}$  in particular, to be a valid marker. There were three studies where no significant relationship was found. In the first, Nugent et al. [41] did not find a significant relationship between  $\dot{V}O_{2\text{peak}}$  and postoperative outcome, though there was an underlying trend. The authors suggested that this outcome may have been influenced by small sample sizes and a reduced tendency to operate on clinically borderline patients in AAA repair as opposed to other surgical indications. The other two instances were in oesophagectomy studies.

Nagamatsu et al.'s [40] later study and Forshaw et al. [37] did not observe a relationship between AT and post-operative complication rate. In the former this was hypothesised to be due to oesophagectomy, with three-field lymphadenectomy, having a cardiopulmonary burden that exceeded the exercise workload determined by AT [40]. While in the latter it was proposed that the small proportion of patients with an AT of  $< 11 \text{ ml}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$  (16%), the non-distinction of cardiopulmonary complications from others such as anastomotic leak and the fact all subjects spent their first night postoperatively on ITU equivalent wards may have masked any underlying trend.

One possibility is that some types of surgery maybe more sensitive to one of these two indicators than the other, that the thresholds for different types of surgery are different (as suggested by Forshaw et al. [37]), and that the level of pre-existing morbidity accepted for surgery may affect the predictive value of the CPET. Thus while we might generally conclude that CPET provides a valid prognostic indicator, this needs to be validated prior to use in specific surgical procedures. The weaker support for AT as a predictor may be a reflection of the uncertainty surrounding the exact physiological significance of this measure.

The data that was used in this review has certain limitations. As with all studies reviewing published results publication bias has the potential to influence our conclusions. However there are other limiting factors specific to this analysis. These include the range of CPET values used as pre-operative indicators, the variability of the defined end point both in terms of time after operation and classification of morbidity and the heterogeneous nature of the subject groups. One study, Older et al. [5], investigated the package of care patients received postoperatively and did not specifically examine the predictive value of CPET data. However it was included in the review as the pre-operative CPET data made up a substantial component of the pre-operative assessment and it demonstrated the increased mortality in the high risk patients even with intensive postoperative support. As a result by demonstrating that patients defined as high risk by AT (a CPET variable) still had a higher mortality despite more intensive care provides further evidence for the ability of CPET to differentiate between levels of postoperative risk.

In Older et al.'s later study [5] elderly patients undergoing intra-abdominal surgery were assigned into one of three classifications of pre and postoperative care, based primarily on their AT. Of the nine cardiopulmonary related deaths, seven came from the most high risk category and two were in the middle category. This demonstrates that using indicators recorded during CPET it is possible to pre-emptively and accurately identify

**Table 2** The ability of  $\dot{V}O_{2peak}$  and anaerobic threshold to predict postoperative cardiopulmonary complications in the studies reviewed. \*Hazard ratios (95% CI) and [SE].

Author	$\dot{V}O_{2peak}/max$				
	Outcome measure	Study result	Summary findings	Anaerobic threshold	
Carlisle [35]	Postoperative survival	Significant predictor	Univariate analysis: $\dot{V}O_{2peak}$ 0.83 (0.74–0.94 [0.05]) $p = 0.002^*$ Multivariate analysis: $\dot{V}_E/\dot{V}CO_2$ : 1.13(1.07–1.19 [0.03]) $p < 0.001^*$ Non-survivors more likely to have a $\dot{V}O_{2peak} < 60\%$ predicted ( $p = 0.04$ ). Non-survivors more likely to have a combined $\dot{V}O_{2peak} < 60\%$ and a $\dot{V}O_2-AT < 50\%$ predicted ( $p < 0.01$ ). Significantly different $\dot{V}O_{2peak}$ between patients who did or did not experience postoperative cardiopulmonary complications ( $p = 0.04$ )	Postoperative survival	Significant predictor Univariate analysis: AT: 0.74 (0.63–0.87 [0.06]) $p < 0.001^*$ Multivariate analysis: AT: 0.84 (0.72–0.98 [0.07]) $p = 0.033^*$ Non-survivors more likely to have a $\dot{V}O_2-AT < 50\%$ predicted ( $p = 0.03$ )
Epstein [36]	Postoperative survival	Significant predictor		Postoperative survival	Significant predictor
Forshaw [37]	Postoperative complications	Significant predictor		Postoperative complications	Not a significant predictor
McCullough [38]	Postoperative complications	Significant predictor	Complications occurred in 16.6% of patients with a peak $\dot{V}O_2 < 15.8 \text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ and in only 2.8% of patients with a peak $\dot{V}O_2 > 15.8 \text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ (univariate odds ratio 6.77 (95% CI 1.14–51.76) $p = 0.002$ ). $\dot{V}O_2$ peak as a continuous variable was a significant predictor of complications (odds ratio of 1.61 per unit decrease (95% CI 1.19–2.18) $p = 0.002$ . Significant difference between the $\dot{V}O_{2max}\cdot\text{m}^{-2}$ of the complications and no complications group ( $p < 0.001$ ). $\dot{V}O_{2max}\cdot\text{m}^{-2}$ was significantly lower in patients with cardiopulmonary complications ( $p < 0.001$ ).	Postoperative complications	Not documented / insufficient analysis AT was not investigated comprehensively but showed a significant trend across the tertiles ( $p < 0.001$ ).
Nagamatsu [39]	Postoperative complications	Significant predictor		Postoperative complications	Significant predictor
Nagamatsu [40]	Postoperative complications	Significant predictor		Postoperative complications	Not a significant predictor
Nugent [41]	Postoperative complications	Not a significant predictor	No significant difference between the $\dot{V}O_{2peak}$ of the complications group (18.6 $\text{ml}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$ ) and the $\dot{V}O_{2peak}$ of the no complications group (21.8 $\text{ml}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$ )	–	Not documented / insufficient analysis



**Table 2** (Continued).

Author	VO <sub>2peak</sub> /max				
	Outcome measure	Study result	Summary findings	Anaerobic threshold	
	Outcome measure	Study result	Summary findings	Outcome measure	
Older [42]	-	Not documented/ insufficient analysis	-	Postoperative survival	Study result
					Summary findings
Older [5]**	-	Not documented/ insufficient analysis	-	-	-

\*\*This study formed patient groups based on AT, type of surgical procedure, ECG changes and V<sub>E</sub>/VO<sub>2</sub>. There was no statistical analysis in the paper of patient outcome based on AT alone though it is strongly suggestive of being a significant predictor.

patients requiring an increased level of postoperative care.  $\dot{V}O_{2max/peak}$  was not used in this study but if this was found to add greater discriminatory power it would further narrow down the high risk patient sub group and so minimise the unwarranted expense of patients unnecessarily receiving more intensive care.

The framework for utilising CPET data with other important predictors of peri-operative risk already exists and provides potential for individualised estimations [43]. However it only currently employs data on a subject's MET (actual peak MET in comparison to their age predicted value). By further understanding the role of the other variables recorded in a CPET and obtaining procedure specific data it may be possible to increase its predictive capacity and improve risk prediction as a whole.

The impact of exercise training on  $\dot{V}O_{2peak}$  in patients has been investigated by several studies. In one study [44] subjects underwent supervised endurance cycle ergometry at 60–100%  $\dot{V}O_{2peak}$  5 days a week for 4–6 weeks prior to their elective operation. Subjects who had an adherence of  $\geq 80\%$  increased their  $\dot{V}O_{2peak}$  by 3.3 ml.kg<sup>-1</sup>.min<sup>-1</sup> (95% CI 1.1–5.4) p = 0.006, representing an increase of 21.5%. This improvement in  $\dot{V}O_{2peak}$  has also been shown using home exercise [45] for 10–12 weeks with weekly telephone monitoring (11% increase in  $\dot{V}O_{2peak}$ , p < 0.05). This was less effective than the first study but as subjects followed a different regime it is not possible to allocate this difference to either the supervision or exercise program followed. Both these studies however demonstrate that in a relatively short pre-surgical period patients are able to significantly elevate their  $\dot{V}O_{2peak}$ . It is yet to be investigated whether this improvement will translate into a lower complication rate from their baseline (pre-fitness regime). However if it did it could represent a powerful method of improving postoperative morbidity and mortality without significant delay to their pre-operative waiting period or adding substantial cost.

There is no firmly identified causal mechanism in the literature which would link a higher  $\dot{V}O_{2peak}$  or AT with an improved risk profile. One possible explanation is that fitter patients are simply more able to mount the sustained increase in oxygen delivery induced by surgery without exceeding their aerobic physiological parameters. A more intriguing hypothesis is that regular exercise can induce a systemic effect similar to ischemic preconditioning which lessens the impact of any imbalance in oxygen delivery and demand associated with surgery, by increasing the body's ability to extract oxygen and better tolerate ischemic conditions. There is both structural and biochemical evidence to support this second possible mechanism. Firstly endurance exercise in humans is known to increase mitochondrial mass [46], which may improve

oxygen utilisation by mitochondria and thus delay the onset of anaerobic respiration and ischemia in the tissues. Secondly animal studies [47] have shown that exercise causes a number of physiological cardiac changes, in particular an increase in the antioxidant capacity and the number of sarcolemmal ATP-sensitive potassium channels. These are proposed to respectively reduce reactive oxygen species damage and cytosolic  $\text{Ca}^{2+}$  overload [47] thereby helping the tissues deal with ischemic conditions. These adaptations may make individuals better able to tolerate periods of anaerobic metabolism peri-operatively.

To gain further evidence to support the use of CPET as a predictor of postoperative morbidity and mortality further studies need to be conducted. These studies need to focus on several key areas, particularly: (i) further investigation of the physiology of AT and the clinical relevance of this phenomenon, (ii) can other variables ( $\text{O}_2$  pulse,  $\dot{V}_E/\dot{V}\text{CO}_2$  etc) independently or combined with  $\dot{V}\text{O}_{2\text{max/peak}}$  and AT provide a more powerful index, and if so would this be able to further narrow down the high risk patients without producing unacceptable type II error, as has been shown in other clinical applications [48]? (iii) Can the risk threshold of the variables be refined for different types of major surgery? (iv) Does increasing a subject's  $\dot{V}\text{O}_{2\text{max}}$  or AT pre-operatively significantly improve their postoperative outcome? (v) What, if any, is the significance of the variables recorded during the recovery phase of the test?

## Acknowledgements

The authors would like to thank Professor Brian Whipp for his comments on this manuscript. The authors are grateful for support from the NIHR Biomedical Research Centre Funding Scheme.

## References

- Matsuoka H, Nishio W, Sakamoto T, Harada H, Tsubota N. Prediction of morbidity after lung resection with risk factors using treadmill exercise test. *European Journal of Cardio-Thoracic Surgery* 2004; **26**: 480–2.
- Bayram AS, Candan T, Gebitekin C. Preoperative maximal exercise oxygen consumption test predicts postoperative pulmonary morbidity following major lung resection. *Respirology* 2007; **12**: 505–10.
- Bolliger CT, Jordan P, Soler M, et al. Exercise capacity as a predictor of postoperative complications in lung resection candidates. *American Journal of Respiratory and Critical Care Medicine* 1995; **151**: 1472–80.
- Mancini DM, Eisen H, Kussmaul W, Mull R, Edmunds LH Jr, Wilson JR. Value of peak exercise oxygen consumption for optimal timing of cardiac transplantation in ambulatory patients with heart failure. *Circulation* 1991; **83**: 778–86.
- Older P, Hall A, Hader R. Cardiopulmonary exercise testing as a screening test for perioperative management of major surgery in the elderly. *Chest* 1999; **116**: 355–62.
- Bylund-Fellenius AC, Walker PM, Elander A, Holm S, Holm J, Schersten T. Energy metabolism in relation to oxygen partial pressure in human skeletal muscle during exercise. *Biochemical Journal* 1981; **200**: 247–5.
- Karlsson J. Pyruvate and lactate ratios in muscle tissue and blood during exercise in man. *Acta Physiologica Scandinavica* 1971; **81**: 455–8.
- Sahlin K, Katz A, Henriksson J. Redox state and lactate accumulation in human skeletal muscle during dynamic exercise. *Biochemical Journal* 1987; **245**: 551–6.
- Wasserman K, Hansen JE, Sue DY, Stringer WW, Whipp BJ. *Principles of Exercise Testing and Interpretation Including Pathophysiology and Clinical Applications*, 4th edn. Philadelphia: Lippincott Williams & Wilkins, 2005.
- Tanaka K, Matsuura Y, Matsuzaka A, Hirakoba K, Kumagai Sun SO, Asano K. A longitudinal assessment of anaerobic threshold and distance-running performance. *Medicine and Science in Sports and Exercise* 1984; **16**: 278–82.
- Zoldatz JA, Sargeant AJ, Emmerich J, Stoklosa J, Zychowski A. Changes in acid-base status of marathon runners during incremental field test. *European Journal of Applied Physiology and Occupational Physiology* 1993; **67**: 71–6.
- Albouuaini K, Eged M, Alahmar A, Wright DJ. Cardiopulmonary exercise testing and its application. *Heart* 2007; **93**: 1285–92.
- Whipp BJ. Carotid bodies and breathing in humans. *Thorax* 1994; **49**: 1081–4.
- Agostoni P, Guazzi M. Evolving role of cardiopulmonary exercise testing in heart failure and cardiac transplantation. In: Weisman IM, Zeballos RJ, eds. *Clinical Exercise Testing*. Basel Karger: Progress in Respiratory Research, 2002: 32: 99–108.
- Beaver WL, Wasserman K, Whipp BJ. A new method for detecting the anaerobic threshold by gas exchange. *Journal of Applied Physiology* 1986; **60**: 2020–7.
- Myers J, Ashley E. Dangerous curves: a perspective on exercise, lactate, and the anaerobic threshold. *Chest* 1997; **111**: 787–95.
- Yeh MP, Gardener RM, Adams TD, Yanowitz FG, Crapo RO. Anaerobic threshold: problems of determination and validation. *Journal of Applied Physiology* 1983; **55**: 1178–86.
- Dennis SC, Noakes TD, Bosch AN. Ventilation and blood lactate increase exponentially during incremental exercise. *Journal of Sports Sciences* 1992; **10**: 437–49.
- Campbell ME, Hughson RL, Green HJ. Continuous increase in blood lactate concentration during different ramp exercise protocols. *Journal of Applied Physiology* 1989; **66**: 1104–7.
- Hughson RL, Weisiger KH, Swanson GD. Blood lactate concentration increase as a continuous function in progressive exercise. *Journal of Applied Physiology* 1987; **62**: 1975–81.
- Gladden LB, Yates JW, Stremel RW, Stamford BA. Gas exchange and lactate anaerobic thresholds: inter- and

- intra-evaluator agreement. *Journal of Applied Physiology* 1985; **58**: 2082–9.
- 22 Paterson DJ, Friedland JS, Bascom DA, et al. Changes in arterial  $K^+$  and ventilation during exercise in normal subjects and subjects with McArdle's Syndrome. *Journal of Physiology* 1990; **429**: 339–48.
  - 23 Henson LC, Ward DS, Whipp BJ. Effect of dopamine on ventilatory response to incremental exercise in man. *Respiration Physiology* 1992; **89**: 209–24.
  - 24 Pandit JJ, Robbins PA. Respiratory effects of breathing high oxygen during incremental exercise in humans. *Advances in Experimental Medicine and Biology* 2002; **499**: 331–6.
  - 25 Busso T, Robbins PA. Evaluation of estimates of alveolar gas exchange by using a tidally ventilated nonhomogenous lung model. *Journal of Applied Physiology* 1997; **82**: 1963–71.
  - 26 Swanson GD. Breath-to-breath considerations for gas exchange kinetics. In: Cerretelli P, Whipp BJ, eds. *Exercise, Bioenergetics and Gas Exchange*. Amsterdam: Elsevier, 1980: 211–22.
  - 27 Pandit JJ, Duncan T, Robbins PA. Total oxygen uptake with two maximal breathing techniques and the tidal volume breathing technique: a physiologic study of pre-oxygenation. *Anesthesiology* 2003; **99**: 841–6.
  - 28 Buchfuhrer MJ, Hansen JE, Robinson TE, Sue DY, Wasserman K, Whipp BJ. Optimising the exercise protocol for cardiopulmonary assessment. *Journal of Applied Physiology* 1983; **55**: 1558–64.
  - 29 Milani RV, Lavie CJ, Mehra MR, Ventura HO. Understanding the basics of cardiopulmonary exercise testing. *Mayo Clinic Proceedings* 2006; **81**: 1603–11.
  - 30 Hansen JE, Sue DY, Wasserman K. Predicted values for clinical exercise testing. *American Review of Respiratory Disease* 1984; **129**: S49–55.
  - 31 American Thoracic Society and the American College of Chest Physicians. ATS/ACCP statement on cardiopulmonary exercise testing. *American Journal of Respiratory and Critical Care Medicine* 2003; **167**: 211–77.
  - 32 Ainsworth BE, Haskell WL, Whitt MC, et al. Compendium of physical activities: an update of activity codes and MET intensities. *Medicine and Science in Sports and Exercise* 2000; **32**: S498–S516.
  - 33 Fleg JL, Pina IL, Balady GJ, et al. Assessment of functional capacity in clinical and research applications: an advisory from the Committee on exercise, rehabilitation, and prevention, Council on clinical cardiology, American Heart Association. *Circulation* 2000; **102**: 1591–7.
  - 34 Beckles MA, Spiro SG, Colice GL, Rudd RM. The physiologic evaluation of patients with lung cancer being considered for resectional surgery. *Chest* 2003; **123**: 105–14.
  - 35 Carlisle J, Swart M. Mid-term survival after abdominal aortic aneurysm surgery predicted by cardiopulmonary exercise testing. *British Journal of Surgery* 2007; **94**: 966–9.
  - 36 Epstein SK, Freeman RB, Khayat A, Unterborn JN, Pratt DS, Kaplan MM. Aerobic capacity is associated with 100-day outcome after hepatic transplantation. *Liver Transplantation* 2004; **10**: 418–24.
  - 37 Forshaw MJ, Strauss DC, Davies AR, et al. Is cardiopulmonary exercise testing a useful test before oesophagectomy? *Annals of Thoracic Surgery* 2008; **85**: 294–9.
  - 38 McCullough PA, Gallagher MJ, deJong AT, et al. Cardiorespiratory fitness and short-term complications after bariatric surgery. *Chest* 2006; **130**: 517–25.
  - 39 Nagamatsu Y, Yamana H, Fujita H, et al. The simultaneous evaluation of preoperative cardiopulmonary functions of oesophageal cancer patients in the analysis of expired gas with exercise testing. *Nippon Kyobu Geka Gakkai Zasshi* 1994; **42**: 2037–40.
  - 40 Nagamatsu Y, Shima I, Yamana H, Fujita H, Shirouzu K, Ishitake T. Preoperative evaluation of cardiopulmonary reserve with the use of expired gas analysis during exercise testing in patients with squamous cell carcinoma of the thoracic esophagus. *Journal of Thoracic and Cardiovascular Surgery* 2001; **121**: 1064–8.
  - 41 Nugent AM, Riley M, Megarry J, O'Reilly MJ, MacMahon J, Lowry R. Cardiopulmonary exercise testing in the pre-operative assessment of patients for repair of abdominal aortic aneurysm. *Irish Journal of Medical Science* 1998; **167**: 238–41.
  - 42 Older P, Smith R, Courtney P, Hone R. Preoperative evaluation of cardiac failure and ischemia in elderly patients by cardiopulmonary exercise testing. *Chest* 1993; **104**: 701–4.
  - 43 Carlisle J. Perioperative risk estimation using survival epidemiology. *Preoperative Association Newsletter*. Feb 2008. [http://www.pre-op.org/Perioperative\\_risk\\_estimation\\_using\\_survival\\_epidemiology.pdf](http://www.pre-op.org/Perioperative_risk_estimation_using_survival_epidemiology.pdf). Last accessed 12 May 2009.
  - 44 Jones LW, Peddle CJ, Eves ND, et al. Effects of presurgical exercise training on cardiorespiratory fitness among patients undergoing thoracic surgery for malignant lung lesions. *Cancer* 2007; **110**: 590–8.
  - 45 Debigare R, Maltais F, Whittom F, Deslauriers J, LeBlanc P. Feasibility and efficacy of home exercise training before lung volume reduction. *Journal of Cardiopulmonary Rehabilitation and Prevention* 1999; **19**: 235–41.
  - 46 Robergs RA, Roberts SO. *Fundamental Principles of Exercise Physiology: for Fitness, Performance, and Health*, 1st edn. Boston: McGraw Hill, 2000.
  - 47 Powers SK, Quindry JC, Kavazis AN. Exercise-induced cardioprotection against myocardial ischemia-reperfusion injury. *Free Radical Biology and Medicine* 2008; **44**: 193–201.
  - 48 Gitt AK, Wasserman K, Kilkowski C, et al. Exercise anaerobic threshold and ventilatory efficiency identify heart failure patients for high risk of early death. *Circulation* 2002; **106**: 3079–84.